

8700200

THE UNITED SHAVES OF ANTERIOA

TO ALL TO WHOM THESE; PRESENTS; SHALL COME:

Nickerson American Plant Breeders, Inc.

Telliereas. There has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF CIGhteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT ETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT, UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Abilene'

In Lestimony Whercot, I have hereunto set my hand and caused the seal of the Plant Variety Exotection Office to be affixed at the City of Washington, D. C. this 30th day of December in the year of our Lord one thousand nine hundred and eighty-eight.

Vielad E. Lyng

Socrotary of Agriculture

Allest

Ternell H. wars

Plant Variety Protection Office Agricultural Marketing Service

The state of the s

				ROVAL EXPIRES 4-30-85				
U.S. DEPARTMENT AGRICULTURAL M.		ED: OMB NO. 0581-0055						
				uired in order to determine protection certificate is to				
APPLICATION FOR PLANT VARI	ETY PROTE	CTION CERTIFICATE	be issued (7 U.S.	C. 2421). Information is				
	s on reverse)		(7 U.S.C. 2426).	until certificate is issued				
1. NAME OF APPLICANT(S)		2. TEMPORARY DESIGNATIO	N 3. VARIETY NA	AME				
Nickerson American Plant Bree	eders Inc.	W81-362-5	Abilene	:				
4. ADDRESS (Street and No. or R.F.D. No., City, Sta.	FOR OFF	ICIAL USE ONLY						
5201 Johnson Drive	1 15 10 10							
Mission, Kansas 66205		303-532-3721 CO	87	00200				
6. GENUS AND SPECIES NAME		(S	IDATE	00200				
	7. FAMILY NAI	6.5	1 1 2	st31,1987				
Triticum estivum	Gramin	eae	LINE TIME	— — — — — — — — — — — — — — — — — — —				
8. KIND NAME	IND NAME							
6. KIND NAME	1_1001							
Hard Red Winter Wheat	ard Red Winter Wheat							
	2=1985							
10. IF THE APPLICANT NAMED IS NOT A "PERSO	AMOUNT	ST 31,1987						
partnership, association, etc.)	₩ s 200) <u>6-0</u>						
Corporation	L DATE							
		:1,1988						
11. IF INCORPORATED, GIVE STATE OF INCORPO		CORPORATION						
13. NAME AND ADDRESS OF APPLICANT REPRES	ENTATIVE(S) II	ANY TO SERVE IN THIS APP	L January	19, 1983				
R.E. Heiner	•		ns or C. Bru					
5201 Johnson Drive	0		econd St.	113				
Mission, KS 66205 (913)384-4940 Routhoud CO 80512								
PHONE Thickdot area code): (303)532-3721								
14 CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED 2 X Exhibit A Origin and Receding History of the Variety (See Section 52 of the Plant Variety Protection 4 of 1)								
b. Exhibit B, Novelty Statement.	a. Exhibit A, Origin and Breeding History of the Variety (See Section 52 of the Plant Variety Protection Act.) b. Exhibit B, Novelty Statement.							
c. Exhibit C, Objective Description of Variety		· · · · · · · · · · · · · · · · · · ·	fice.)	*				
d. X Exhibit D, Additional Description of Varie			•					
c. X Exhibit E, Statement of the Basis of Appli	cant's Ownership	. f. Exhibit F. Qua	lity & Agron	omic Data 💮				
15. DOES THE APPLICANTIS) SPECIFY THAT SEED SEED? (See Section 83(a) of the Plant Variety Pro	OF THIS VARIE	y my		—				
16. DOES THE APPLICANT(S) SPECIFY THAT THIS	SEED? (See Section 83(a) of the Plant Variety Protection Act.) Yes fil "Yes," answer DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE 17. IF "YES" TO ITEM 16,							
LIMITED AS TO NUMBER OF GENERATIONS?		BEYOND BRESDER S	יפוז	_				
18. DID THE APPLICANT(S) PREVIOUSLY FILE F	OR PROTECTI	X Foundation	<u>KX</u> Registered	KX Certified				
PILE PILE APPLICATION THE VIOLOGET FILE P	-ON FROTECTIO	DN OF THE VARIETY IN THE	0.5.7	Yes (If "Yes," give date)				
				No				
19. HAS THE VARIETY BEEN RELEASED, OFFER	ED FOR SALE,	OR MARKETED IN THE U.S. O	R OTHER COUNT	RIES ? Yes (If "Yes," give names				
				of countries and dates)				
			N N	No				
20. The applicant(s) declare(s) that a viable sampl	e of basic seeds	of this variety will be furnished	ed with the applica	tion and will be re-				
plenished upon request in accordance with suc The undersigned applicant(s) is (are) the owne distinct, uniform, and stable as required in Sec	r(s) of this sexu	ally reproduced novel plant v	ariety, and believe he provisions of Se	(s) that the variety is ection 42 of the Plant				
Variety Protection Act. Applicant(s) is (are) informed that false repres	sentation berein	can leonardize protection and	l result in nenaltie	5				
	as Possasso.							
SIGNATURE OF APPLICANT	DATE							
SIGNATURE OF APPLICANT			DATE 7-20					
Robert Burn.			I ' '					
SIGNATURE OF APPLICANT SIGNATURE OF APPLICANT De 11 e			I ' '					
Robert Bur.			7-20					

EXHIBIT A

ORIGIN AND BREEDING HISTORY OF ABILENE

Abilene was derived in 1983 as a single F7 head-row reselection from the segregating motherline W81-362. W81-362 was an F3 derived line from the cross OK11252/W76-1226, which had superior performance in yield trials but was recycled in head-rows due to excessive levels of segregation.

Abilene was tested under the experimental line number W81-362-5 in preliminary and replicated yield trials at numerous locations in the Hard Red Winter Wheat Region of the Great Plains from 1984-1986. It was entered in the 1986 Southern Regional Performance Nursery (SRPN) and was the highest yielding line over locations. It is entered in the 1987 SRPN, and official state and private trials in Texas, Oklahoma, Kansas, Colorado, Nebraska, Missouri, New Mexico and South Dakota for further testing.

In 1984 .005 of an acre derived from a single F7 head was grown as a preliminary Breeder Seed Increase. In 1985 450 pounds of Breeder Seed was produced in Berthoud, Colorado.

Abilene is uniform and stable. Less than .5% of the plants were rogued from the foundation fields in 1986. Approximately 95% of the rogued variant plants were 3 to 12 centimeters taller than Abilene. Less than .5% total variant plants may be encountered in subsequent generations.

EXHIBIT B

NOVELTY STATEMENT

Abilene is most similar to the hard red winter wheat Trailblazer. However, it can be distinguished by the following morphological charateristics:

- -Abilene has a blue-green plant color at boot stage. Trailblazer has a green plant color at boot stage.
- -Abilene and Trailblazer both have acuminate type beak lengths. However, Abilene's beak length is significantly longer than Trailblazer's (see statistical data following page).
- -Abilene has a glume length classified as medium.
 Trailblazer has a glume length classified as long, (see statistical data following page).
- -Both Abilene and Trailblazer differ significantly in plant height (see statistical data following page).

TABLE FOR GLUME LENGTH ABILENE VERSUS TRAILBLAZER

SOURCE	<u>DF</u>		<u>ss</u>	<u>MS</u>
TOTAL	49		29.642	
VAR	1	the second second	22.984	22.98400
ERROR	48		6.658	0.13870

F-TEST = 165.713** CV = 0.495 LSD(5%) = 0.042

MEANS FOR EACH VARIETY

ABILENE = 7.96 mm TRAILBLAZER = 9.32 mm

** The difference in means of glume length are significantly different at the 1% probability level.

ANOVA TABLE FOR BEAK LENGTH ABILENE VERSUS TRAILBLAZER

SOURCE	DF	<u>SS</u>	<u>MS</u>
TOTAL	49	238.020	151 27050
VAR	<u>. </u>	151.380	151.37959
ERROR	48	86.641	1.80502

F-TEST = 83.866** CV = 3.926 LSD(5%) = 0.152

MEANS FOR EACH VARIETY

ABILENE = 7.17 mm TRAILBLAZER = 3.69 mm

** The difference in means of beak length are significantly different at the 1% probability level.

ANOVA TABLE FOR PLANT HEIGHT ABILENE-MESA-TRAILBLAZER

SOURCE	<u>DF</u>	<u>SS</u>	<u>MS</u>
TOTAL	98	10931.344	
VAR	2	671.889	335.94460
ERROR	96	10259.455	106.86932

F-TEST = 6.287** CV = .207 LSD(5%) = .886

MEANS FOR EACH VARIETY

ABILENE = 69.24 cm MESA = 67.12 cm TRAILBLAZER = 73.39 cm

**The difference in means of plant height are significantly different at the 1% probability level.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT, GRAIN & SEED DIVISION BELTSVILLE, MARYLAND 20708 EXHIBIT C (Wheat)

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

NAME OF APPLICANTIS	FOR OFFICIAL USE ONLY
Nickerson American Plant Breeders Inc.	PYPO NUMBER
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	8700200
5201 Johnson Drive	VARIETY NAME OR TEMPORARY DESIGNATION
Mission, Kansas 66205	
Place the appropriate number that describes the varietal character of this v Place a zero in first box (e-8- 0 8 9 or 0 9) when number is either 9	
1. KIND:	
1 1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 = POLISH	6 = POULARD 7 = CLUB
1	SOFT 3 = OTHER (Specify) HARD
2 1 = WHITE 2 = RED 3 = OTHER (Specify)	
3. SEASON - NUMBER OF DAYS FROM planting To	
2 3 4 FIRST FLOWERING 2 4	0 LAST FLOWERING
4. MATURITY (50% Flowering): EQUAL TO TRAILBLAZER	
<u>, </u>	1 = ARTHUR 2 = SCOUT 3 = CHRIS
NO. OF DAYS LATER THAN	4=LEMHI 5=NUGAINES 6=LEEDS 7=Trailblaze
5. PLANT HEIGHT (From soil level to top of head):	
0 6 9 CM. HIGH	
CM. TALLER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
CM SHORTER THAN	1 = ARTHUR 2 = SCOUT 4 = LEMHI 5 = NUGAINES 6 = LEEDS 7 = Trailblaze
6. PLANT COLOR AT BOOTING (See reverse): 7. ANT	HER COLOR:
3 1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN 1	= YELLOW 2 = PURPLE
8. STEM:	
1 Anthocyanin: 1 = ABSENT 2 = PRESENT - 2	axy bloom: 1 = ABSENT 2 = PRESENT
2 Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT 1 In	tternodes: 1 = HOLLOW 2 = SOLID
0 4 NO. OF NODES (Originating from node above ground)	CM. INTERNODE LENGTH BETWEEN FLAG LEAF AND LEAF BELOW
9. AURICLES:	
2 Anthocyanin: 1 = ABSENT 2 = PRESENT 2 H	airiness: I = ABSENT 2 = PRESENT
19. LEAF:	•
Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify): 2 F	lag leaf: 1 = NOT TWISTED 2 = TWISTED
Hairs of first leaf sheath: 1 = ASSENT 2 = PRESENT 2	axy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT
1 4 MM. LEAF WIDTH (First leaf below flag leaf)	3 CM. LEAF LENGTH (First leaf below fing leaf):
200 470 C 17 G 17 G 17 (Favors of 3.70 may be used)	

Abil	ene '		8700200
	HEAD:		<u> </u>
	Density: 1 = LAX 2 = DENSE 3=Middense	Shape: 1 = TAPERING 4 = OTHER (Spe	
	Awnedness: 1 = AWNLESS 2 = APICALLY AWNLETED 3	= AWNLETED ' 4 = AWNED	
	1=WHITE 2=YELLOW 3=PINK 4=	RED	
	Color at maturity: 5 = BROWN 6 = BLACK 7 = OTHE	R (Specify):	
[O. 1 CM. LENGTH	1 1 mm. wioth	
12.	GLUMES AT MATURITY:		
2	-	1 Width: 1 = NARROW (C. 3 = WIDE (CA. 4	
2	Shoulder 1 = WANTING 2 = OBLIQUE 3 = ROUNDED shape: 4 = SQUARE 5 = ELEVATED 6 = APICULATE	3 Beak: T=OBTUSE	2 = ACUTE 3 = ACUMINATE
13.	COLEOPTILE COLOR:	14. SEEDLING ANTHOCYANI	N:
	- 7		
	I = WHITE 2 = RED 3 = PURPLE	2 1 = ABSENT 2 = P	RESENT
15.	JUYENILE PLANT GROWTH HABIT:		
2	! = PROSTRATE 2 = SEMI-ERECT 3 = EREC	τ	
16.	SEED:		
1	Shape: 1 = OVATE 2 = OVAL 3 = ELLIPTICAL	1 Cheek: 1 = ROUNDED	2 = ANGULAR
2	Brush: 1 = SHORT 2 = midlong 3 = Long	Brush: = NOT COLL	ARED 2 = COLLARED
[-	Phenol reaction 1 = IVORY • 2 = FAWN 3 = LT. BROWN (See instructions): 4 = BROWN 5 = BLACK	·	
[7	£	
	3 Color: 1 = WHITE 2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)	
	5 .0 MM. LENGTH 3.3 MM. WIDTH	3 0 GM. PER 1000 SEI	EDS
17.	SEED CREASE:		
	Vidth: 1 = 60% OR LESS OF KERNEL 'WINOKA'	1 7 1	ESS OF KERNEL 'SCOUT'
	2 = 80% OR LESS OF KERNEL 'CHRIS'		ESS OF KERNEL 'CHRIS'
·	3 = NEARLY AS WIDE AS KERNEL 'LEMHI'		ESS OF KERNEL 'LEMHI'
. 18.	DISEASE: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 3=MC	derately Susceptible	4=Moderately Resistant
	STEM RUST	and the first of the contract	A LOOSE SMUT
4	[(Races) [7/4] (Races)field_race	S (Racea)	
	POWDERY MILDEW 0 BUNT		Borne Mosaic Virus
19.	INSECT: (0 = Not Tested, 1 = Susceptible, 2 = Resistant) 3=MC	derately Susceptible	4= <u>Mod</u> erately Resistant
	SAWFLY 0 APHID (Bydy.)	0 GREEN BUG	O CEREAL LEAF BEETLE
	OTHER (Specify) HESSIAN FLY	1 GP 0 A	0 B 0 C
	RACES: (0 E	0 F 0 G
30	INDICATE WHICH VARIETY HAST OF ASE, V RESENOTES THAT	SIBULTIED:	
20.	CHARACTER NAME OF VARIETY	CHARACTER	NAME OF VARIETY
	Plant tillering Trailblazer	Seed size	Trailblazer
	Leaf size Trailblazer	Seed shape	Trailblazer
	. , , ,	Coleoptile elongation	Trailblazer
		Seedling pigmentation	Trailblazer
		<u> </u>	TRATIDIAZEL
	INSTRU	CTIONS	

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

EXHIBIT D

ADDITIONAL BOTANICAL DESCRIPTION OF ABILENE

Abilene is a hard red winter wheat bred and developed by Nickerson American Plant Breeders Inc. It was tested as experimental number W81-362-5.

Abilene is a very high yielding, medium maturity, short to intermediate semidwarf wheat with very good levels of winterhardiness and good milling and baking properties. Abilene has consistently been in the top yield group throughout much of the HRWW region.

In addition to its high yield potential Abilene has been identified as having; excellent test weight patterns, good drought response, very good to excellent straw strength, very good protection to the prevalent races of leaf rust and speckled leaf blotch, excellent protection to soil borne mosaic virus, fair to good protection against tan spot and wheat spindle streak mosaic virus. Abilene is susceptible to Hessian fly and powdery mildew.

Juvenile growth habit is semi-erect. Plant color at boot is blue-green with an erect, twisted flag leaf. Head shape is strap to tapering, middense, awned and head color is white at maturity. Auricle anthocyanin and auricle hairs are present. Glumes are midlong and narrow with oblique shoulders and acuminate beaks. Seed shape is ovate with rounded cheeks. Seed crease is narrow and shallow.

Abilene's primary area of adaptation has tentatively been identified as northern two-thirds of the HRWW region (Blackwell, Oklahoma, and north), plus the panhandle regions of Oklahoma and Texas. Abilene was the highest yielding line over locations in the 1986 Southern Regional Performance Nursery (SRPN) indicating that it has a very broad area of adaptation in the central Great Plains.

EXHIBIT E

STATEMENT OF THE BASIS OF APPLICANT'S OWNERSHIP

Nickerson American Plant Breeders Inc. is the applicant for protection in this case being:

- a) The incorporated business (registered in Delaware) for and within which regular employees have bred the named variety.
- b) The proprietory owner and intending commercial user of the variety.

EXHIBIT F. QUALITY AND AGRONOMIC DATA

Quality Da	ata .					•	•	•	• •	•	•	.page	1.
Agronomic	Data	and	Patho:	logical	Traits		•		•			. page	2 2.

								AGRIFRO	SEEDS							:		
YEAR	1986	-			,*	·	HARD RE	RED WINTER	R WHEAT	QUALITY	<u>≻</u>		•				PAGE	- 1
				WHEA	WHEATFLO	UR QUALITY	-ITY		•	-	BAKING	GUALITY	<u>}</u>					
YEAR	SAMPLE		TEST WT.	FROT	FLR	FLR	FLR ASH	MIX	ABS.	MIX	CHAR	LOAF	GRN	CRUMB	[]g	MILL	BAKE	TOTAL
AB.	ABILENE		1b/Bu	14%mb	×	14%mb	14%mb	ŭ	×	nie.	E.	000	ŭ	œ	ır	1 1 1 1 1 1		
94	HW81-362-5	85	6.09	13.5	71.6	12.5	0.000	មាល	63.0	ម ម		1000+	000	0) [100) (((((((((((((((((((A-08	0 (1)
תור נסיני	HW91-362-5		5.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00		69.1	12.5	0.430	v (\	64.0	กตุ น่อ	\ K	0 to 0	(ە ر	4 n	~ 0	7 14 7 15 7 15 7 15 7 15 7 15 7 15 7 15 7 15	67 - D	164-E
(O)	HW81-362-5		61.5	12.9	69.6	12.4	0.407	۰.	64.0) ()	- 1\	.096)	. o)) (0)) <u>an</u>) (n - a)	(A) (A) (A) (A) (A)	169-B
85	HW81-362-5		e0.e	12.6	68.6	11.3	0.393	۲.	64.0	(V)	۲.	880	٨.	0)	o)	2-62	90-E	153-0
90	W81-362-5		0 0 0		64.5	11.9	0.411	φ	62.0	4	0)	1000+	. 0)	0)	0)	J-T2	91 0) 0)	159-0
96	MST-362-2		59.7		62.0	11.3	0.487	វ្រា	62.0	თ (რ	ø	980	Û	0)	0)	64-D	84-E	1430-0
9	W81-362-5	GI.	61.9	12.3	65.5	10.9	0.408	۸.	62.0	79 10	a)	900	۸.	o)	တ	2-0-L	西 - 四 6	153-C
	AVERAGE	Ή	59.9	12.7	67.6	11.7	0.424	Ψ	63.1	60	og .	983	K	5)	0)	74-C	82-B	159-0
94	NOT MEN	So	57.5	!	69.3	10.5	0.000	4	60.0	, e	0)	960	6	1)))	62-D)-8½	140-C
9)	NUMATOR	브	59.7		67.9	11.5	0.509	N	65.0	61 10	0)	810	2	0)	1.	2-92	78-0	154-C
(0 (0	NEWTON	쏫	52.1	12.6	66.0	11.6	0.431	ln	eo.o	0.0	t	840	រភ	w	۷.	50-F	67-D	Q-955
(c 2)	NEMTON	BB	2.95	ଧ• ୯୯	99	12.2	0.378	r.	64.0	ტ დ	ø	1000	0)	Ú)	ထ	73-0	91-A	164-B
ဗ	NEUTON	チ	54.8	13.6	66.7	12.2	0.40€	မ	62.0	0) (n	۷	930	0)	ص	ou	67-D	96-18	122-C
9	NEMLON	2	54.5	12.0	63.7	10.9	0.000	K	63.0	4.0	Ø	006	t~	۲.	0)	€.0-D	86-B	146-C
98	NEWTON	춪	52,7		e0.e	ν.	0.420	വ	63.0	e) : () :	<i>ا</i> د	280	o) i	0) 1	O) :	47-F	28-C	125-0
9	NEETON	E9	e0°3	11.4	ນ ພິ	a)	0.407	o O	64.0	ν.	œ	000	Φ)	0)	0)	6.2-D	81-B	143-C
	AVERAGE	Й	56.4	12.2	66.0	11.1	0.425	φ	62.€	(O	o)	873		o)	0)	64-D	80-B	144-C
			 		Í 					11 11 14 14 14 14 14	***************************************							1
Ď	GRADES: 6	A-EXCELLENT	ENT	B- 6000		C-ACCEPTABLE	ABLE	(A)	D-QUESTIONABLE	BLE	ır.	F-UNACCEPTABLE	TABLE			. •		
R=RA		9-10-EXCELLENT	EZ.	8=1300D		ACCEPT	ABLE	5-6=Ω∪	ESTIONA	ELE E	1-4=∟	INACCEF	TABLE					

RELATIVE RANKINGS OF ABILENE AND 17 OTHER HRWW VARIETIES FOR VARIOUS AGRONOMIC AND PATHOLOGICAL TRAITS

VARIETY	TEST WEIGHT	SHATTERING POTENTIAL	ANTHESIS	PHYSIOLOGICAL MATURITY	COLEOPTILE EXPRESSION	HEIGHT	STRAW STRENGTH	WINTERHARDINESS	LEAF RUST	STEM RUST	SOIL BORNE MOSAIC VIRUS	SPINDLE STREAK MOSAIC VIRUS	SEPTORIA TRITICI	TAN SPOT	POWDERY MILDEW	HESSIAN FLY
Abilene Trailblazer Mesa Siouxland Brule Rocky Tam 107 Arkan Vona Tam 105 Newton Wings Hawk Wrangler Mustang Thunderbird Victory	34346465665263446	65668647434533366	553567245345555555	6545562344463353		-443657343444335 4	25256734335653434	35532434737555634	3 4 3 7 7 8 4 7 8 8 7 8 2 3	64425451685544525	23386262862833312	44375254994527661	23535553546467734	5 2 7 4 7 5 6 5 5 5 7 5 4 7 4 7 4	7862841245544776 5	-8847288248848664 8
Stallion	4	6	5	. 4	5	3	1	6	6	7	3	6	3	6	6	4

Mean Score 4.5 5.1 6.7 4.9 6.2 5.3 5.1 6.4 6.7 4.3 4.9 5.7 4.4 5.3 5.5 5.8 Std. Dev. 1.3 1.5 3.9 1.9 4.1 3.9 3.2 3.8 3.3 1.9 3.9 3.6 1.5 1.4 2.2 2.4

The rankings in the table above are based on a scale of 1-9, where 1 and 9 represent the following extremes for the respective traits.

·	1	9
Test Weight	high	low
Shattering Potential Maturity	low early	high late
Coleoptile Expression Straw Strength	short strong	long weak
Winterkill All disease & insect ratings	hardy resistant	tender susceptible